



Heterogeneous Microbial Populations: using flow cytometric data for building dynamic distributed models

Lencastre Fernandes, Rita; Carlquist, M.; Lundin, L.; Heins, Anna-Lena; Dutta, Abhishek; Nopens, I.; Johansen, Søren J.; Jensen, Anker Degn; Eliasson Lantz, Anna; Gernaey, Krist

Publication date:
2011

[Link back to DTU Orbit](#)

Citation (APA):

Lencastre Fernandes, R., Carlquist, M., Lundin, L., Heins, A-L., Dutta, A., Nopens, I., Johansen, S. J., Jensen, A. D., Eliasson Lantz, A., & Gernaey, K. (2011). *Heterogeneous Microbial Populations: using flow cytometric data for building dynamic distributed models*. Abstract from 8th European Congress of Chemical Engineering, Berlin, Germany.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Heterogeneous Microbial Populations: using flow cytometric data for building dynamic distributed models

R. Lencastre Fernandes, M. Carlquist, Technical University of Denmark, Kgs. Lyngby, Denmark; L. Lundin, University of Copenhagen, Denmark, A.-L. Heins, Technical University of Denmark, Kgs. Lyngby, Denmark; A. Dutta, I. Nopens, Ghent University, Belgium; A.D. Jensen, A. Eliasson Lantz, K.V. Gernaey, Technical University of Denmark, Kgs. Lyngby, Denmark

Traditionally, microbial populations have been considered homogeneous in studies of fermentation processes. However, research has shown that a typical microbial population in a fermentor is heterogeneous [1-3].

Phenotypic heterogeneity arises as a result of the variability inherent to the metabolic processes within single cells. Two dominant cell variables responsible for differential gene expression are cell cycle and cell ageing [4]. Indeed, cells at different phases in the cell cycle, or with different ages, have been observed to respond differently to stress conditions [1].

Although the number of experimental methods available for single-cell analysis has boomed [5, 6], the knowledge acquired by such experimental studies has not yet been integrated into a generally accepted modeling framework able to account for distributed properties within a cell population [3].

In this work, focus was set on experimentally studying, as well as modeling, the dynamics of phenotypic heterogeneous populations of *Saccharomyces cerevisiae* during batch cultivations. Besides the common monitored variables (e.g. optical density, glucose, ethanol), single-cell total protein content and DNA content were measured by flow cytometry during the different phases of batch cultivations. Aiming at establishing a population balance model (PBM) which describes the dynamic behavior of the yeast population (including the relative contribution of different subpopulations), a systematic analysis of the flow cytometric data was performed, and mathematical descriptions for the budding initiation and cell division rates as functions of the available substrate concentration are proposed.

1. Avery SV. Microbial cell individuality and the underlying sources of heterogeneity. *Nat Rev Microbiol* 2006; 4:577-587.
2. Enfors SO, Jahic M, Rozkov A, Xu B, Hecker M, Jürgen B *et al.* Physiological responses to mixing in large scale bioreactors. *J Biotechnol* 2001; 85:175-185.
3. Müller S, Harms H, Bley T. Origin and analysis of microbial population heterogeneity in bioprocesses. *Curr Opin Biotechnol* 2010; 21:100-113.
4. Sumner ER, Avery SV. Phenotypic heterogeneity: differential stress resistance among individual cells of the yeast *Saccharomyces cerevisiae*. *Microbiology* 2002M; 148:345-351.
5. Schmid A, Kortmann H, Dittrich PS, Blank LM. Chemical and biological single cell analysis. *Curr Opin Biotechnol* 2010; 21:12-20.
6. Lencastre Fernandes R, Nierychlo M, Lundin L, Pedersen AE, Puentes Tellez PE, Dutta A *et al.* Experimental methods and modeling techniques for description of cell population heterogeneity. Submitted to *Biotechnology Advances* 2011